

# PRESCRIPCIÓN PARA PIOJOS DE LA CABEZA



INDICADO PARA PACIENTES  
MAYORES DE 6 MESES DE EDAD

## Spinosad

Topical Suspension 0.9%

[www.spinosadrx.com](http://www.spinosadrx.com)

Favor leer información importante de seguridad al reverso

# ¡ELIMINE LOS PIJOS CON SPINOSAD SUSPENSIÓN TÓPICA 0.9%!

**Se estima que se producen entre 6 y 12 millones de infestaciones de piojos al año en los niños de edad escolar.<sup>2</sup>**

Spinosad Suspensión Tópica 0.9% es un tratamiento contra piojos de prescripción indicado para utilizar en adultos y niños a partir de los 6 meses de edad. Spinosad Suspensión Tópica 0.9% se debe utilizar en el contexto de un programa para el control general de la pediculosis. Spinosad Suspensión Tópica 0.9% se aplica en forma tópica en el cuero cabelludo y cabello para eliminar la infestación de piojos.

Los piojos son diminutos insectos sin alas que no vuelan ni saltan, pero trepan, como medio principal de transportación. La principal forma de contagio de piojos es por el contacto de cabeza a cabeza.

El ciclo de vida del piojo tiene 3 etapas: huevo (liendre) que está adherida al cabello, ninfa, y adjunto (ver imagen a continuación). Los piojos y sus huevos en general se encuentran en la base del cabello y cerca del cuero cabelludo. El piojo adulto es aproximadamente del tamaño de una semilla de sésamo.<sup>3</sup>



**Ver la Información completa para el paciente sobre la prescripción adjunta. Spinosad Suspensión Tópica 0.9% está disponible solo con prescripción médica.**

# Resultados clínicos comprobados para una condición de comparación directa.


Recientemente, evidencia clínica ha demostrado que los piojos han desarrollado resistencia a dos tipos de insecticidas, la piretrina y los piretroides, incluida la permetrina, que es el ingrediente activo utilizado en un tratamiento sin receta comúnmente utilizado para las infestaciones de piojos (Nix® [permetrina 1%]).<sup>4</sup>

Spinosad Suspensión Tópica 0.9% es el primer tratamiento de receta aprobado por la Administración de Drogas y Alimentos de EE. UU. (FDA) que, en dos estudios clínicos de comparación directa, demostró que:


- Spinosad Suspensión Tópica 0.9% es más efectivo para eliminar infestaciones de piojos que la permetrina 1% (también conocida como Nix®).<sup>5</sup> (Ver los gráficos a continuación)
- NO es necesario usar el peine fino para piojos con Spinosad Suspensión Tópica 0.9%.
- FUE necesario usar el peine fino para piojos con Nix® (permetrina 1%).

Variable Principal: La proporción de pacientes sin piojos (sin piojos vivos, adultos o ninfas) 14 días después del último tratamiento con Spinosad Suspensión Tópica 0.9% sin peinar versus permetrina con peinar.

## Study 1: Treatment Success



## Study 2: Treatment Success



# Spinosad

## Topical Suspension 0.9%

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Spinosad Topical Suspension 0.9% safely and effectively. See full prescribing information for Spinosad Topical Suspension 0.9%.

Spinosad Topical Suspension 0.9%, for topical use.  
Initial U.S. Approval: 2011

#### RECENT MAJOR CHANGES

Indications and Usage (1)

12/2014

#### INDICATIONS AND USAGE

Spinosad Topical Suspension 0.9% is a pediculicide indicated for the topical treatment of head lice infestations in patients six (6) months of age and older. (1.1)

#### DOSAGE AND ADMINISTRATION

- For topical use only. Not for oral, ophthalmic, or intravaginal use. (2)
  - Shake bottle well. (2)
  - Apply product to dry scalp and hair using only the amount needed to cover the scalp and hair. (2)
  - Rinse off with warm water after 10 minutes. (2)
- Repeat treatment only if live lice are seen 7 days after first treatment. (2)

#### DOSAGE FORMS AND STRENGTHS

Suspension: 9 mg of spinosad per gram of Spinosad Topical Suspension 0.9% in 120 mL bottles (3)

#### CONTRAINDICATIONS

None. (4)

#### WARNINGS AND PRECAUTIONS

- Benzyl alcohol toxicity: Not recommended in infants below the age of 6 months; potential for increased systemic absorption. (5.1)

#### ADVERSE REACTIONS

Most common adverse events (>1%) were application site erythema and ocular erythema. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact ParaPRO, LLC at 1-855-628-7622 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

See 17 for PATIENT COUNSELING INFORMATION and FDA approved patient labeling.  
Revised: 12/2014

### FULL PRESCRIBING INFORMATION: CONTENTS\*

- 1 INDICATIONS AND USAGE
  - 1.1 INDICATION
  - 1.2 ADJUNCTIVE MEASURES
- 2 DOSAGE AND ADMINISTRATION
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
  - 5.1 BENZYL ALCOHOL TOXICITY
- 6 ADVERSE REACTIONS
  - 6.1 CLINICAL STUDIES EXPERIENCE
- 8 USE IN SPECIFIC POPULATIONS
  - 8.1 PREGNANCY

8.3 NURSING MOTHERS

8.4 PEDIATRIC USE

8.5 GERIATRIC USE

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 MECHANISM OF ACTION

12.2 PHARMACODYNAMICS

12.3 PHARMACOKINETICS

## **13 NONCLINICAL TOXICOLOGY**

13.1 SPINOSAD: CARCINOGENESIS, MUTAGENESIS, IMPAIRMENT OF FERTILITY

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

16.1 HOW SUPPLIED

16.2 STORAGE AND HANDLING

## **17 PATIENT COUNSELING/INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

# **FULL PRESCRIBING INFORMATION**

Spinosad Topical Suspension 0.9%

## **1 INDICATIONS AND USAGE**

### **1.1 Indication**

Spinosad Topical Suspension 0.9% is indicated for the topical treatment of head lice infestation in patients six (6) months of age and older.

### **1.2 Adjunctive Measures**

Spinosad Topical Suspension 0.9% should be used in the context of an overall lice management program:

- Wash (in hot water) or dry-clean all recently worn clothing, hats, used bedding and towels.
- Wash personal care items such as combs, brushes and hair clips in hot water
- A fine-tooth comb or special nit comb may be used to remove dead lice and nits.

## **2 DOSAGE AND ADMINISTRATION**

For topical use only. Spinosad Topical Suspension 0.9% is not for oral, ophthalmic, or intravaginal use.

Shake bottle well. Apply sufficient Spinosad Topical Suspension 0.9% to cover dry scalp, then apply to dry hair. Depending on hair length, apply up to 120 mL (one bottle) to adequately cover scalp and hair. Leave on for 10 minutes, then thoroughly rinse off Spinosad Topical Suspension 0.9% with warm water. If live lice are seen 7 days after the first treatment, a second treatment should be applied. Avoid contact with eyes.

## **3 DOSAGE FORMS AND STRENGTHS**

0.9%, viscous, slightly opaque, light orange-colored suspension in 120 mL bottles.

## **4 CONTRAINDICATIONS**

None.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Benzyl Alcohol Toxicity**

Spinosad Topical Suspension 0.9% contains benzyl alcohol and is not recommended for use in neonates and infants below the age of 6 months. Systemic exposure to benzyl alcohol has been associated with serious adverse reactions and death in neonates and low birth-weight infants [See *Use in Specific Populations* (8.4)].

## **6 ADVERSE REACTIONS**

### **6.1 Clinical Studies Experience**

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in clinical practice.

Spinosad Topical Suspension 0.9% was studied in two randomized, active-controlled trials (N=552) in subjects with head lice; the results are presented in Table 1.

**Table 1: Selected Adverse Events Occurring in at least 1% of Subjects**

Signs	Spinosad (N=552)	Permethrin 1% (N=457)
Application site erythema	17 (3%)	31(7%)
Ocular erythema	12 (2%)	15(3%)
Application site irritation	5 (1%)	7(2%)

Other less common reactions (less than 1% but more than 0.1%) were application site dryness, application site exfoliation, alopecia, and dry skin.

Systemic safety was not assessed in pediatric subjects under 6 months of age as laboratory parameters were not monitored in these controlled studies.

**8 USE IN SPECIFIC POPULATIONS**

**8.1 Pregnancy**

Pregnancy Category B.

There are no adequate and well-controlled studies with Spinosad Topical Suspension 0.9% in pregnant women. Reproduction studies conducted in rats and rabbits were negative for teratogenic effects. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

No comparisons of animal exposure with human exposure are provided in this labeling due to the low systemic exposure noted in the clinical pharmacokinetic study [see *Clinical Pharmacology* (12.3)] which did not allow for the determination of human AUC values that could be used for this calculation.

Systemic embryofetal development studies were conducted in rats and rabbits. Oral doses of 10, 50 and 200 mg/kg/day spinosad were administered during the period of organogenesis (gestational days 6 - 15) to pregnant female rats. No teratogenic effects were noted at any dose. Maternal toxicity occurred at 200 mg/kg/day. Oral doses of 2.5, 10, and 50 mg/kg/day spinosad were administered during the period of organogenesis (gestational days 7 - 19) to pregnant female rabbits. No teratogenic effects were noted at any dose. Maternal toxicity occurred at 50 mg/kg/day.

A two-generation dietary reproduction study was conducted in rats. Oral doses of 3, 10, and 100 mg/kg/day spinosad were administered to male and female rats from 10 - 12 weeks prior to mating and throughout mating, parturition, and lactation. No reproductive/developmental toxicity was noted at doses up to 10 mg/kg/day. In the presence of maternal toxicity, increased dystocia in parturition, decreased gestation survival, decreased litter size, decreased pup body weight, and decreased neonatal survival occurred at a dose of 100 mg/kg/day.

**8.3 Nursing Mothers**

Spinosad, the active ingredient in Spinosad Topical Suspension 0.9% is not systemically absorbed; and therefore, will not be present in human milk. However, Spinosad Topical Suspension 0.9% contains benzyl alcohol, which may be systemically absorbed through the skin, and the amount of benzyl alcohol excreted in human milk with use of Spinosad Topical Suspension 0.9% is unknown. Caution should be exercised when Spinosad Topical Suspension 0.9% is administered to a lactating woman. A lactating woman may choose to pump and discard breast milk for 8 hours (5 half-lives of benzyl alcohol) after use to avoid infant ingestion of benzyl alcohol.

**8.4 Pediatric Use**

The safety and effectiveness of Spinosad Topical Suspension 0.9% have been established in pediatric patients 6 months of age and older with active head lice infestation [see *Clinical Pharmacology* (12.3) and *Clinical Studies* (14)].

Safety in pediatric patients below the age of 6 months has not been established.

Spinosad Topical Suspension 0.9% is not recommended in pediatric patients below the age of 6 months because of the potential for increased systemic absorption due to a high ratio of skin surface area to body mass and the potential for an immature skin barrier.

Spinosad Topical Suspension 0.9% contains benzyl alcohol which has been associated with serious adverse reactions and death in neonates and low birth-weight infants. The "gasping syndrome" (characterized by central nervous system depression, metabolic acidosis, gasping respirations, and high levels of benzyl alcohol and its metabolites found in the blood and urine) has been associated with benzyl alcohol dosages >99 mg/kg/day in neonates and low birth-weight infants when administered intravenously. Additional symptoms may include gradual neurological

deterioration, seizures, intracranial hemorrhage, hematologic abnormalities, skin breakdown, hepatic and renal failure, hypotension, bradycardia, and cardiovascular collapse.

The minimum amount of benzyl alcohol at which toxicity may occur is not known. Premature and low birth-weight infants, as well as patients receiving high dosages, may be more likely to develop toxicity [see *Warning and Precautions* (5.1)].

### 8.5 Geriatric Use


Clinical studies of Spinosad Topical Suspension 0.9% did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

## 10 OVERDOSAGE

If oral ingestion occurs, seek medical advice immediately.

## 11 DESCRIPTION

Spinosad Topical Suspension 0.9%, is a slightly opaque, light orange colored, viscous topical suspension. Spinosad, the active ingredient, is derived from the fermentation of a soil actinomycete bacterium, *Saccharopolyspora spinosa*. Spinosad is a mixture of spinosyn A and spinosyn D in a ratio of approximately 5 to 1 (spinosyn A to spinosyn D).



**Spinosyn A:** The chemical name is: 1H-as-Indaceno[3,2-d]oxacyclododecin-7,15-dione, 2-[(6-deoxy-2,3,4-tri-O-methyl- $\alpha$ -L-mannopyranosyl)oxy]-13-[[2R,5S,6R)-5-(dimethylamino)tetrahydro-6-methyl-2H-pyran-2-yl]oxy]-9-ethyl-2,3,3a,5a,5b,6,9,10,11,12,13,14,16a,16b-tetradecahydro-14-methyl-, (2R,3aS,5aR,5bS,9S,13S,14R,16aS,16bR)-

**Spinosyn D:** The chemical name is: 1H-as-Indaceno[3,2-d]oxacyclododecin-7,15-dione, 2-[(6-deoxy-2,3,4-tri-O-methyl- $\alpha$ -L-mannopyranosyl)oxy]-13-[[2R,5S,6R)-5-(dimethylamino)tetrahydro-6-methyl-2H-pyran-2-yl]oxy]-9-ethyl-2,3,3a,5a,5b,6,9,10,11,12,13,14,16a,16b-tetradecahydro-4,14-dimethyl-, (2S,3aSR,5aS,5bS,9S,13S,14R,16aS,16bS)-

Spinosad Topical Suspension 0.9% contains 9 mg spinosad per gram in a viscous, slightly opaque, light orange colored vehicle consisting of Water, Isopropyl Alcohol, Benzyl Alcohol, Hexylene Glycol, Propylene Glycol, Cetearyl Alcohol, Stearalkonium Chloride, Cetareth-20, Hydroxyethyl Cellulose, Butylated Hydroxytoluene, FD&C Yellow #6.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Spinosad causes neuronal excitation in insects. After periods of hyperexcitation, lice become paralyzed and die.

### 12.2 Pharmacodynamics

The pharmacodynamics of Spinosad Topical Suspension 0.9% has not been studied.

### 12.3 Pharmacokinetics

An open-label, single-center trial was conducted over a period of seven days to determine the pharmacokinetic profile of spinosad 1.8% in pediatric subjects with head lice infestation. Fourteen (14) subjects, 4 - 15 years of age, with head lice were enrolled into the trial. All subjects applied a single topical (scalp) treatment

treatment of spinosad 1.8% for 10 minutes, after which the test article was washed off, and subjects underwent plasma sampling. Results demonstrated that spinosad was below the limit of quantitation (3ng/mL) in all samples. Plasma concentration of benzyl alcohol was not determined in these subjects. An open-label, two-center trial was conducted over a period of 23 days to determine the pharmacokinetic profile of spinosad 0.9% and the ingredient benzyl alcohol in pediatric subjects with a head lice infestation. Twenty-six (26) subjects between 6 months to 4 years of age were enrolled into the study per protocol. All subjects applied a single topical (scalp) treatment of spinosad 0.9% for 10 minutes, after which the test article was washed off, and subjects underwent plasma sampling over a 12-hour period. Plasma spinosad concentrations were below the limit of quantitation (3 ng/mL) in all samples.

Benzyl alcohol was quantifiable (above 1 µg/mL) in a total of 8 plasma samples in 6 out of 26 subjects (25%): four out of 12 subjects in the 6 months to < 2 years age group and two out of 14 subjects in the 2 to 4 years age group. The highest observed concentration was 2.37 µg/mL. Benzyl alcohol concentrations at 12 hours post-treatment were below limit of quantification (1 µg/mL) for all subjects.

### 13 NONCLINICAL TOXICOLOGY

#### 13.1 Carcinogenesis, Mutagenesis, Impairment of fertility

In an oral (diet) mouse carcinogenicity study, spinosad was administered to CD-1 mice at doses of 0.0025, 0.008, and 0.036% in the diet (approximately 3.4, 11.4, and 50.9 mg/kg/day for males and 4.2, 13.8, and 67.0 mg/kg/day for females) for 18 months. No treatment-related tumors were noted in the mouse carcinogenicity study up to the highest doses evaluated in this study of 50.9 mg/kg/day in male mice and 13.8 mg/kg/day in female mice. Female mice treated with a dose of 67.0 mg/kg/day were not evaluated in this study due to high mortality.

In an oral (diet) rat carcinogenicity study, spinosad was administered to Fischer 344 rats at doses of 0.005, 0.02, 0.05, and 0.1% in the diet (approximately 2.4, 9.5, 24.1 and 49.4 mg/kg/day for males and 3.0, 12.0, 30.1 and 62.8 mg/kg/day for females) for 24 months. No treatment-related tumors were noted in the rat carcinogenicity study in male or female rats up to the highest doses evaluated in this study of 24.1 mg/kg/day in male rats and 30.1 mg/kg/day in female rats. Rats in the highest dose group in this study were not evaluated due to high mortality.

Spinosad demonstrated no evidence of mutagenic or clastogenic potential based on the results of four *in vitro* genotoxicity tests (Ames assay, mouse lymphoma L5178Y assay, Chinese hamster ovary cell chromosome aberration assay, and rat hepatocyte unscheduled DNA synthesis assay) and one *in vivo* genotoxicity test (mouse bone marrow micronucleus assay).

Oral administration of spinosad (in diet) to rats, throughout mating, gestation, parturition and lactation, demonstrated no effects on growth, fertility or reproduction, at doses up to 10 mg/kg/day [see *Pregnancy* (8.1)].

### 14 CLINICAL STUDIES

Two multicenter, randomized, investigator-blind, active-controlled studies were conducted in 1,038 subjects 6 months of age and older with head lice infestation. A total of 552 subjects were treated with Spinosad Topical Suspension 0.9%. For the evaluation of efficacy, the youngest subject from each household was considered to be the primary subject of the household, and other members in the household were enrolled in the study as secondary subjects, and evaluated for all safety parameters.

In Study 1, 91 primary subjects were randomized to Spinosad Topical Suspension 0.9%, and 89 primary subjects were randomized to permethrin 1%. In Study 2, 83 and 84 primary subjects were randomized to Spinosad Topical Suspension 0.9% and permethrin 1%, respectively.

In both studies, all subjects who were treated on Day 0 returned for efficacy evaluation at Day 7. Subjects with live lice present at Day 7 received a second treatment. Subjects who were lice free on Day 7 were to return on Day 14 for evaluation. Subjects with live lice and who received a second treatment were to return on Days 14 and 21.

Efficacy was assessed as the proportion of primary subjects who were free of live lice 14 days after the final treatment. Table 2 contains the proportion of primary subjects who were free of live lice in each of the two trials.



# Pon fin al ciclo del piojo

Table 2. Proportion of Subjects Free of Live Lice 14 days After Last Treatment

Study 1		Study 2	
Spinosad Topical Suspension 0.9% N=91	permethrin 1% N=89	Spinosad Topical Suspension 0.9% N=83	permethrin 1% N=84
77 (84.6%)	40 (44.9%)	72 (86.7%)	36 (42.9%)

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**16.1 How Supplied**

Spinosad Topical Suspension 0.9% is a slightly opaque, light orange colored, viscous liquid, supplied in 4 oz. (120 mL) high density polyethylene (HDPE) bottles. NDC 28595-570-04

**16.2 Storage and Handling**

- Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F).
- Keep out of reach of children

**17 PATIENT COUNSELING/INFORMATION**

[See FDA-approved patient labeling (Patient Information)]

The patient should be instructed as follows:

- Shake bottle well immediately prior to use.
- Do not swallow.
- Use Spinosad Topical Suspension 0.9% only on dry scalp and dry scalp hair.
- Repeat treatment only if live lice are seen seven days after first treatment.
- Avoid contact with eyes. If Spinosad Topical Suspension 0.9% gets in or near the eyes, rinse thoroughly with water.
- Wash hands after applying Spinosad Topical Suspension 0.9%.
- Use Spinosad Topical Suspension 0.9% on children only under direct supervision of an adult.

## Patient Information

### Spinosad Topical Suspension 0.9%

**Important: For use on scalp hair and scalp only. Do not get Spinosad Topical Suspension 0.9% in your eyes, mouth, or vagina.**

Read the Patient Information that comes with Spinosad Topical Suspension 0.9% before you start using it and each time you get a refill. There may be new information. This leaflet does not take the place of talking to your healthcare provider about your medical condition or treatment.

**What is Spinosad Topical Suspension 0.9%?**

Spinosad Topical Suspension 0.9% is a prescription medicine used to get rid of lice in scalp hair of children and adults.

It is not known if Spinosad Topical Suspension 0.9% is safe for children under 6 months of age or in people over age 65.

Once Spinosad Topical Suspension 0.9% is washed off, a fine-tooth comb may be used to remove treated lice and nits from the hair and scalp, but combing is not required. All personal items exposed to the hair or lice should be washed in hot water or dry-cleaned. See “How do I stop the spread of lice?” at the end of this leaflet.

**What should I tell my healthcare provider before I use Spinosad Topical Suspension 0.9%?**

**Tell your healthcare provider about all of your medical conditions, or the medical conditions of your child, including if you or your child:**

- have any skin conditions or sensitivities
- are pregnant or planning to become pregnant. It is not known if Spinosad Topical Suspension 0.9% can harm your unborn baby.
- are breastfeeding. Talk to your healthcare provider if you are breastfeeding.

**How should I use Spinosad Topical Suspension 0.9%?**

- Use Spinosad Topical Suspension 0.9% exactly as prescribed. Your healthcare provider will prescribe the treatment that is right for you. Do not change your treatment unless you talk to your healthcare provider.

- Use Spinosad Topical Suspension 0.9% in one or two treatments that are one week apart. If live lice are seen one week (7 days) after you first used Spinosad Topical Suspension 0.9%, you will need to use Spinosad Topical Suspension 0.9% again.
- Shake bottle well right before use.
- Use Spinosad Topical Suspension 0.9% when your hair is dry. Do not wet your hair before applying Spinosad Topical Suspension 0.9%.
- It is important to use enough Spinosad Topical Suspension 0.9% to coat completely every single louse and to leave it on your scalp for the full 10 minutes. See the detailed Patient Instructions for Use at the end of this leaflet.
- Because you need to completely cover all of the lice with Spinosad Topical Suspension 0.9%, you may need help in applying Spinosad Topical Suspension 0.9% to your scalp and hair. Make sure that you and anyone who helps you apply Spinosad Topical Suspension 0.9% reads and understands this leaflet and the Patient Instructions for Use.
- Children will need an adult to apply Spinosad Topical Suspension 0.9% for them.
- Do not swallow Spinosad Topical Suspension 0.9%. If swallowed, call your healthcare provider right away.
- Do not get into eyes. If Spinosad Topical Suspension 0.9% gets in the eye, flush with water right away.
- Wash your hands after you apply Spinosad Topical Suspension 0.9%.

### **What are the possible side effects of Spinosad Topical Suspension 0.9%?**

People using Spinosad Topical Suspension 0.9% may have skin or eye:

- Redness
- Irritation

If skin or eye irritation happens, rinse with water right away, then call your healthcare provider or go to the emergency department.

These are not all the side effects of Spinosad Topical Suspension 0.9%. For more information, ask your healthcare provider.

Call your doctor for medical advice about side effects. You may report side effects to FDA at **1-800-FDA-1088**.

### **How should I store Spinosad Topical Suspension 0.9%?**

- Store Spinosad Topical Suspension 0.9% in a dry place at room temperature, 20°C to 25°C (68°F to 77°F).

### **Keep Spinosad Topical Suspension 0.9% and all medicines out of the reach of children.**

### **What are the ingredients in Spinosad Topical Suspension 0.9%?**

Active ingredient: spinosad

Inactive ingredients: water, isopropyl alcohol, benzyl alcohol, hexylene glycol, propylene glycol, cetearyl alcohol, stearylalkonium chloride, cetareth-20, hydroxyethyl cellulose, butylated hydroxytoluene, FD&C yellow #6

### **General Information about Spinosad Topical Suspension 0.9%**


Medicines are sometimes prescribed for conditions other than those described in the patient information leaflets. Do not use Spinosad Topical Suspension 0.9% for any condition for which it was not prescribed by your healthcare provider. Do not give Spinosad Topical Suspension 0.9% to other people, even if they have the same symptoms as you. It may harm them.

This leaflet summarizes the most important information about Spinosad Topical Suspension 0.9%. If you would like more information, talk to your healthcare provider. You can also ask your healthcare provider for information about Spinosad Topical Suspension 0.9% that is written for healthcare professionals.

## **Patient Instructions for Use**

Completely cover the scalp with Spinosad Topical Suspension 0.9%. Lice and nits live near the scalp. Use as much product as needed to completely cover the scalp first, and then apply outwards towards the ends of the hair.

For very thick, medium length hair or long hair, an entire bottle (120 mL) of Spinosad Topical Suspension 0.9% may be needed to cover the scalp and hair. Less Spinosad Topical Suspension 0.9% may be needed for shorter, thinner hair.



## Step 1

- Shake Spinosad Topical Suspension 0.9% bottle well right before use.



## Step 2

- Cover your face and eyes with a towel and keep your eyes closed tightly.
- Apply Spinosad Topical Suspension 0.9% directly to **dry** hair.
- Completely cover the scalp first, and then apply outwards towards the ends of the hair.
- If not enough Spinosad Topical Suspension 0.9% is used, some lice may escape treatment. It is important to use enough Spinosad Topical Suspension 0.9% to cover your entire scalp and all scalp hair.



## Step 3

- Allow Spinosad Topical Suspension 0.9% to stay on your hair for 10 minutes. Use a timer or clock and start timing after you have completely covered your hair and scalp with Spinosad Topical Suspension 0.9%.
- Continue to keep eyes covered to prevent dripping into your eyes.
- After 10 minutes, completely rinse Spinosad Topical Suspension 0.9% from your hair and scalp with warm water.
- You or anyone who helps you apply Spinosad Topical Suspension 0.9% should wash hands after application.
- It is okay to shampoo your hair any time after the treatment.



One week (7 days) after your first treatment, if live lice are seen, repeat the steps above.

## How do I stop the spread of lice?

To help prevent the spread of lice from one person to another, here are some steps you can take:

- Avoid direct head-to-head contact with anyone known to have live, crawling lice.
- Do not share combs, brushes, hats, scarves, bandannas, ribbons, barrettes, hair bands, towels, helmets, or other hair-related personal items with anyone else, whether they have lice or not.
- Avoid sleepovers and slumber parties during lice outbreaks. Lice can live in bedding, pillows, and carpets that have recently been used by someone with lice.
- After finishing treatment with lice medicine, check everyone in your family for lice after one week. Be sure to talk to your healthcare provider about treatments for those who have lice.
- Machine-wash any bedding and clothing used by anyone having lice. Machine-wash at high temperatures (150°F) and tumble in a hot dryer for 20 minutes.

**This patient leaflet has been approved by the U.S. Food and Drug Administration.**

Distributed by: Allegis Pharmaceuticals, LLC, Canton, MS 39046  
SPN-PI-001

**Pon fin al ciclo del piojo**

**Data sugiere que los pacientes pueden tratar las infestaciones de piojos en forma independiente 5 veces antes de consultar con el proveedor de atención médicos.<sup>6</sup>**

**Si sospecha que tiene piojos, consulte con su proveedor de atención médicos sobre...**



**Eficacia comprobada sin necesidad de peine fino**


**Información importante de seguridad**

Spinosad Suspensión Tópica 0.9% contiene alcohol bencílico y su uso no es recomendado en neonatos y niños menores de 6 meses de edad. La exposición sistémica al alcohol bencílico se ha asociado con diferentes reacciones adversas y muerte en neonatos y niños de bajo peso al nacer, cuando se administra en forma intravenosa. Los eventos adversos más comunes fueron: enrojecimiento en lugar de la aplicación (3%) y enrojecimiento ocular (2%) irritación del sitio de aplicación (1%).

# Tan fácil como 1, 2, 3...

## Instrucciones de uso:

Cubra completamente cuero cabelludo con Spinosad Suspensión Tópica 0.9%. Los piojos y las liendres viven cerca del cuero cabelludo. Use la cantidad de producto que sea necesaria para cubrir completamente el cuero cabelludo primero y luego aplique hacia afuera, hacia las puntas del cabello. Para cabellos muy gruesos, de largo medio o largos puede ser necesario utilizar un frasco entero (120 mL) de Spinosad Suspensión Tópica 0.9% para cubrir el cuero cabelludo y el cabello completamente. Para cabello más fino y menos largo, se puede necesitar menos Spinosad Suspensión Tópica 0.9%.



### Paso 1

- Agite el frasco energicamente antes de usar.



### Paso 2

- Cubra el rostro y los ojos con una toalla y mantenga los ojos bien cerrados.
- Aplique Spinosad Suspensión Tópica 0.9% directamente en el cabello seco.
- Cubra completamente el cuero cabelludo primero y luego aplique hacia afuera, hacia las puntas del cabello.
- Si no utiliza suficiente Spinosad Suspensión Tópica 0.9%, algunos piojos pueden escapar el tratamiento. Es importante utilizar suficiente Spinosad Suspensión Tópica 0.9% para cubrir todo el cuero cabelludo y todo el cabello.



### Paso 3

- Deje Spinosad Suspensión Tópica 0.9% en el cabello y el cuero cabelludo durante 10 minutos. Utilice un reloj y comience a medir el tiempo después de cubrir completamente el cabello y cuero cabelludo con Spinosad Suspensión Tópica 0.9%.
- Continúe con los ojos cubiertos para evitar que le caigan gotas del producto en los ojos.
- Después de 10 minutos, enjuague completamente Spinosad Suspensión Tópica 0.9% del cabello y el cuero cabelludo con agua tibia.
- Usted y la persona que le haya ayudado a aplicar Spinosad Suspensión Tópica 0.9% deberán lavarse las manos después de la aplicación.
- Se puede lavar el cabello con champú inmediatamente después del tratamiento.



Una semana (7 días) después del primer tratamiento, si se encuentran piojos vivos, repita los pasos anteriores.

**Importante: Para uso tópico solo sobre el cuero cabelludo y el cabello. No deje que Spinosad Suspensión Tópica 0.9% tenga contacto con los ojos, boca o vagina.**

Consulte la Información para el Paciente adjunta y con su médico para obtener información más completa sobre Spinosad Suspensión Tópica 0.9%.

# Pon fin al ciclo del piojo

## Indicación

Spinosad Suspensión Tópica 0.9% es un producto para el tratamiento tópico de la pediculosis por infestación de piojos en pacientes mayores de seis (6) meses de edad.

## Medidas Adicionales

Spinosad Suspensión Tópica 0.9% se debe utilizar en el contexto de un programa para el control general de la pediculosis:

- Lave (en agua caliente) o limpie en seco toda la ropa, sombreros, ropa de cama y toallas utilizadas recientemente
- Lave todos los elementos de cuidado personal tales como peines, cepillos, y sujetadores de cabello en agua caliente
- Se puede utilizar un peine fino, o un peine especial para sacar liendres, para eliminar los piojos y liendres muertos.

## Información importante sobre seguridad

Spinosad Suspensión Tópica 0.9% contiene alcohol bencílico y su uso no es recomendado en neonatos y niños menores de 6 meses de edad. La exposición sistémica al alcohol bencílico se ha asociado con diferentes reacciones adversas y muerte en neonatos y niños de bajo peso al nacer, cuando se administra en forma intravenosa. Los eventos adversos más comunes fueron: enrojecimiento en lugar de la aplicación (3%) y enrojecimiento ocular (2%) irritación del sitio de aplicación (1%).

**Para ayudar a prevenir la propagación de piojos de una persona a otra, a continuación, se incluyen algunas medidas que se pueden tomar:**

- Evite el contacto cabeza a cabeza con alguna persona que se sepa que está plagada de piojos vivos.
- No comparta artículos personales relacionados con el cabello.
- Evite pijamadas y fiestas en las que los niños se quedan a dormir con amigos durante brotes de piojos.
- Después de terminar el tratamiento contra piojos, revise todos los miembros de la familia para verificar que no tengan piojos después de una semana. Asegúrese de hablar con su proveedor de atención médica sobre los tratamientos que se pueden aplicar a los que tengan piojos.
- Lave a máquina la ropa de cama y vestimenta que hayan usado las personas infestadas.

Consulte la Información para el Paciente adjunta con su médico para obtener información más completa sobre Spinosad Suspensión Tópica 0.9%.

Se recomienda denunciar efectos secundarios negativos de las medicinas de prescripción a la FDA. Visite la página [www.fda.gov/medwatch](http://www.fda.gov/medwatch), o comuníquese al **1-800-FDA-1088**.

# Spinosad

## Topical Suspension 0.9%

1. Spinosad Topical Suspension 0.9% [package insert]
2. <https://www.cdc.gov/parasites/lice/head/epi.html>
3. Frankowsky et al. Pediatrics 2010
4. KJ Gellatly, S Krim, DJ Palenchar, K Shepherd, KS Yoon, CJ Rhodes, SH Lee, JM Clark, Journal of Medical Entomology, Expansion of the Knockdown Resistance Frequency Map for Human Head Lice (Phthiraptera: Pediculidae) in the United States Using Quantitative Sequencing, 2016.
5. Stough et al. Pediatrics 2009
6. West DP. Am J Manage Care 2004